Journal of Elasticity manuscript No.
(will be inserted by the editor)

Translation of Michael Sadowsky's paper "The differential equations of the MÖBIUS band"*

Denis F. Hinz · **Eliot Fried**

arXiv:1408.2747v2 [math.HO] 31 Aug 2014

Received: date / Accepted: date

Abstract This article is a translation of Michael Sadowsky's original paper "Die Differentialgleichungen des MÖBIUSschen Bandes." in Jahresbericht der Deutschen Mathematiker-Vereinigung **39** (2. Abt. Heft 5/8, Jahresversammlung vom 16. bis 23. September), 49–51 (1929), which is a short version of his paper "Theorie der elastisch biegsamen undehnbaren Bänder mit Anwendungen auf das MÖBIUS'sche Band" in 3. internationaler Kongress für technische Mechanik, Stockholm, 1930.

Keywords Möbius band · Energy functional · Bending energy

Translation of the original paper

In a previously completed work, Sadowsky [1] provided a proof for the *existence* of a developable MÖBIUS band and provided a variational context for the underlying geometric problem. In the present work, the problem of determining the equilibrium shape of a MÖBIUS band formed from an elastic strip is treated as a static mechanical problem.

Let P denote an arbitrary point on the midline of the MÖBIUS band, let s denote the arclength of the mid-line, let $\mathbf{t}$, $\mathbf{n}$, and $\mathbf{b}$ denote the accompanying vector triad at P, and let K and W denote the curvature and twist of the midline at P. Considering

* Citations of this translation should refer also to Sadowsky's original paper, as cited in the Abstract.

Denis F. Hinz
Mathematical Soft Matter Unit
Okinawa Institute of Science and Technology
Okinawa, Japan 904-0495
E-mail: dfhinz@gmail.com

Eliot Fried
Mathematical Soft Matter Unit
Okinawa Institute of Science and Technology,
Okinawa, Japan 904-0495
E-mail: eliot.fried@oist.jp

a cut perpendicular to the midline at P, the stresses along the line of the cut may be reduced to a force $\mathfrak{K}$ and a moment $\mathfrak{M}$ both acting at P.

The representations of $\mathfrak{K}$ and $\mathfrak{M}$ relative to the triad $\mathbf{t}$, $\mathbf{n}$, and $\mathbf{b}$ shall be denoted as

$$\mathfrak{K} = T\mathbf{t} + N\mathbf{n} + B\mathbf{b}, \tag{1}$$

and

$$\mathfrak{M} = \mathfrak{T}\mathbf{t} + \mathfrak{N}\mathbf{n} + \mathfrak{B}\mathbf{b}. \tag{2}$$

From a mechanical perspective, the band is defined through the following three requirements:

1. The band is an object with distinguished midline.
2. The vector triad $\mathbf{t}$, $\mathbf{n}$, and $\mathbf{b}$ accompanying the midline is material.
3. The midline is inextensible.

(The second requirement results from the band being part of the rectifying surface of its midline).

Applying a virtual deformation to the band and invoking the Frenet–Serret formulas results in

$$d\delta\vartheta = (\mathbf{t}\delta W + \mathbf{b}\delta K)ds, \tag{3}$$

and in

$$\delta ds = 0, \tag{4}$$

where $\delta\vartheta$ is the virtual rotation of the vector triad at P.

The following considerations hold only for infinitesimally narrow bands.

The virtual work of the internal forces is[1]

$$\delta A_i = -\int_{s_1}^{s_2} (\mathfrak{T}\delta W + \mathfrak{B}\delta K)ds. \tag{5}$$

The mean curvature H of the band surface at P is[1]

$$H = \frac{K^2 + W^2}{2K}. \tag{6}$$

The elastic potential E of the band is proportional to the square of the mean curvature (granted the presumption that the band surface is a developable surface)

$$E = A\frac{(K^2 + W^2)^2}{K^2}. \tag{7}$$

Since $\delta A_i = -\int_{s_1}^{s_2} \delta E ds$ and, in view of (5) and (7), it follows that

$$\mathfrak{T} = \frac{\partial E}{\partial W} = A\frac{4W(K^2 + W^2)}{K^2} \tag{8}$$

[1] HAMEL, Über die Mechanik der Drähte und Seile, Sitzungsberichte der Berliner Mathematischen Gesellschaft, XXV, 1925/26. Also HAMEL Die Axiome der Mechanik, Handbuch der Physik, herausgegeben von H. GEIGER und KARL SCHEEL, Band V. In the latter work further references may be found.

and

$$\mathfrak{B} = \frac{\partial E}{\partial K} = A\frac{2(K^4 - W^4)}{K^3}, \tag{9}$$

where A is a positive material constant. The six components appearing in (1) and (2) satisfy six known *equilibrium equations*.[1] Using (8) and (9), those equations read

$$\begin{cases} T = A\left(C - \dfrac{(K^2+W^2)^2}{K^2}\right), \\ N = -\dfrac{A}{K}\dfrac{d}{ds}\dfrac{(K^2+W^2)^2}{K^2}, \\ B = \dfrac{2AW}{K^3}(K^2+W^2)^2 + 4A\dfrac{d}{ds}\left(\dfrac{1}{K}\dfrac{d}{ds}\dfrac{W(K^2+W^2)}{K^2}\right), \\ \mathfrak{N} = \dfrac{4A}{K}\dfrac{d}{ds}\dfrac{W(K^2+W^2)}{K^2}, \end{cases} \tag{10}$$

and

$$\begin{cases} KT + \dfrac{dN}{ds} - WB = 0, \\ WN + \dfrac{dB}{ds} = 0, \end{cases} \tag{11}$$

where C is an arbitrary constant of integration.

If the expressions (10) were to be used in (11), two differential equations for K and W, respectively, would result. The integration of these equations would yield the natural midline of the band. Equations (8), (9), (10), and (11) shall be referred to as the equations of the MÖBIUS band.

The midline of the band possesses a singular point X that can be found in the following way: a MÖBIUS band possesses a symmetry axis defined such that it is congruent with itself after a rotation of 180° about that axis. The symmetry axis intersects the band at two points such that the axis coincides with the binormal $\mathbf{b}$ at one of these points; this is the point X.

Let φ denote the angle between the rectilinear generators of the band through the point P and $\mathbf{b}$; then[1]

$$\tan\varphi = \frac{W}{K}. \tag{12}$$

For the point X,

$$\mathfrak{B} = 0 \tag{13}$$

due to symmetry.

To arrive at an additional conclusion, consider an *experiment with a band model*: this shows that

$$\lim_{P\to X} \varphi \neq 0, \tag{14}$$

$$\lim_{P\to X} \mathfrak{T} \neq 0. \tag{15}$$

The experiment corresponding to (14) consists of observing a band; for (15), one needs to cut the boundaries of a band at X to infer the moment $\mathfrak{T}$ from the twist

of the band in the weakened cross-section. From (13), (14), (15), and the governing equations (8) and (9) of the band it transpires that

$$\lim_{P\to X} K \neq 0, \quad \lim_{P\to X} W \neq 0, \quad \text{and} \quad \lim_{P\to X} \varphi = 45^\circ. \tag{16}$$

The last limit in (16) points to the following peculiar fact:

A MÖBIUS *band consists of a planar, right triangle. The curved, analytic portion of the band connects continuously with the two legs of the right triangle; it connects with continuous tangential plane, but with discontinuous curvature.*[2]

References

1. M. Sadowsky, Ein elementarer Beweis für die Existenz eines abwickelbaren Möbiusschen Bandes und die Zurückführung des geometrischen Problems auf ein Variationsproblem, Sitzungsberichte der Preussischen Akademie der Wissenschaften, physikalisch-mathematische Klasse **22**, 412–415 (1930)

[2] For an interpretation of the last section of Sadowsky's original paper including figures generated from recent numerical simulations, refer to D. F. Hinz and E. Fried, Translation and interpretation of Michael Sadowsky's paper "Theory of elastically bendable inextensible bands with applications to the MÖBIUS band", submitted to *Journal of Elasticity* (2014).